AU-127

B.Sc. Part—III Semester—V Examination MATHEMATICS (OLD) (UPTO SUMMER-2018) (Analysis)

Paper-IX

Time: Three Hours1 [Maximum Marks: 60] **N.B.**:— (1) Question No. 1 is compulsory. (2) Attempt ONE question from each Unit. 1. Choose the correct alternatives:— A bounded function f is R-integrable on [a, b] if its: (a) upper and lower integrals are equal. (b) upper and lower integrals are not equal. (c) L(P, f) = U(P, f). (d) None of the above. (ii) An improper integral $\int_{a}^{b} \frac{dx}{(b-x)^{p}}$ converges if: 1 (b) p < 1(a) p > 1(d) $p \le 1$ (c) $p \ge 1$ (iii) If f(z) = u + iv be analytic in a region D of z-plane then: 1 (a) u is harmonic but v is not (b) v is harmonic but u is not (c) both u & v are harmonic. (d) neither u is harmonic nor v is harmonic. (iv) A bilinear transformation w = f(z) having only one fixed point is: (a) elliptic (b) loxodromic (c) hyperbolic (d) parabolic VOX--35303 (Contd.)

Ī

(Contd.)

(v) Let A be a subset of a metric space X. Then A is open if:

	(a) For every $x \in A$, there is $r \ge 0$ such that $N_1(x) \subset A$.				
	(b)	(b) each point of A is its interior point.			
	(c)	for every $x \in A$, there is a neighborhood of x that lies entirely in A.			
	(d)	all of these.			
(vi)	Let	$B = \{x/0 \le x \le 1 \text{ and } x \text{ is rational }$	num	ber} be a subset of metric space R. Then: 1	
	(a)	B is closed in R			
	(b)	B is open in R			
	(c)	B is not open in R			
	(d)	None of these.			
(vii)	The	invariant points of the transformat	ion v	$v = \frac{3z - 4}{z - 1} \text{ are} :$	
	(a)	1 ± i	(b)	2, 2	
	(c)	2 ± i	(d)	-2, -2	
(viii)	Set	of Real numbers R is:		1	
	(a)	countable	(b)	uncountable	
	(c)	finite	(d)	None of these.	
(ix)	A n	eighborhood of a point in a metric	spac	e is:	
	(a)	an open set	(b)	a closed set	
	(c)	neither open nor closed	(d)	both open and closed	
(x)	A tr	ransformation $w = f(z) = e^{i\pi/4} \cdot z$ is a	:	1	
	(a)	rotation	(b)	magnification	
	(c)	translation	(d)	inversion	
		UN	IT	·I	
(a)				[a, b] is integrable on [a, b] iff for each $\ensuremath{\varepsilon} \ge 0$	
		partition P of [a, b] such that $U(P, f) - L(P, f) \le \epsilon$.			
(b)	lf f i	is a function defined by $f(x) = x$ on	0, 2],	then show that f is integrable in Riemann sense	
	ove	$f[0, 2] \text{ and } \int_{0}^{2} f(x) dx = z.$		5	

2

VOX -35303

2.

www.sgbauonline.com

- 3. (p) Let f(x), $g(x) \in C$, $a \le x \le \infty$ and $0 \le f(x) \le g(x)$ for every $x \ge a$. Then prove that :
 - (i) $\int_{0}^{\infty} g(x)dx$ converges $\Rightarrow \int_{0}^{\infty} f(x)dx$ converges
 - (ii) $\int_{a}^{\infty} f(x) dx = \infty \Rightarrow \int_{a}^{\infty} g(x) dx = \infty.$
 - (q) Prove that:
 - (i) $\int_{1}^{\infty} \frac{\sin x}{x^2} dx \text{ converges absolutely.}$
 - (ii) $\int_{2}^{\infty} \frac{x^3 dx}{\sqrt{x^2 + 1}}$ diverges.
 - (iii) $\int_{2}^{\infty} \frac{x^{2} dx}{\sqrt{x^{7} + 1}}$ converges.

UNIT-II

- 4. (a) Prove that the function sin z is analytic and find its derivative.
 - (b) Prove that the Cauchy-Riemann equations in polar form can be written as $u_r \frac{1}{r}v_o$ and

 $v_r = -\frac{1}{r}u_o$, where w = f(z) = u(x, y) + iv(x, y) is a function defined in a region D of Z-plane.

- 5. (p) If w = f(z) = u + iv be analytic in a region D then prove that the families of curves $u(x, y) C_1$ and $v(x, y) = C_2$ form an orthogonal system of curves, where C_1 and C_2 are arbitrary constants.
 - (q) Find the analytic function whose real part is $e^{-x}(x \sin y y \cos y)$.

UNIT—III

6. (a) Prove that every bilinear transformation with a single non-infinite fixed point α can be put in the normal form:

$$\frac{1}{w-\alpha} = \frac{1}{z-\alpha} + K$$
, where K is constant.

VOX—35303 3 (Contd.)

www.sgbauonline.com

(b)	Find the region D' of w-plane into which the region D in Z-plane bounded by $x =$	0.					
	$y = 0$, $x \pm y = 1$ is transformed by the mapping $w = e^{i\pi/4} \cdot z$.	5					
(p)	Show that cross ratio remains invariant under a bilinear transformation.	5					
(q)	Find the fixed points of the bilinear transformation $w = \frac{(2+i)z}{i+z} = \frac{2}{i}$. Write its normal form as	nd					
	show that it is Ioxodromic.	5					
UNIT—IV							
(a)	Define:						
	(i) interior point of a set						
	(ii) limit point of a set.	2					
(b)	Prove that every neighborhood is an open set.	3					
(c)	Let (X, d) be a metric space. Then show that d ₁ defined by :						
	$d_1(x,y) = \frac{d(x,y)}{1 + d(x,y)}$						
	is also a metric on X.	5					
(p)	Let $Y \subset X$. Then prove that a subset A of Y is open relative to Y iff A $Y \cap G$ for some open subset G of X.	en 6					
(q)	If (X, d) is a metric space and $A \subset X$, then prove that A is closed iff A contains its boundary	îy. 4					
	UNITV						
(a)	Let $K \subset Y \subset X$. Then prove that K is compact relative to X iff K is compact relative to	Y. 5					
(b)	Let X, Y be metric spaces and $f: X \rightarrow Y$. Prove that f is continuous iff $f(\overline{A}) \subset \overline{F(A)}$ for everywhere A of X.	ry 5					
(p)	Prove that a mapping f of a metric space X into a metric space Y is continuous on X if $f^{-1}(V)$ is open in X for every open set V in Y.	iff 5					
(q)	Prove that every totally bounded metric space is bounded.	5					

VOX 35303

10.

11.

4

775